

Drugs of Abuse

DRUG/FAMILY	AVAILABILITY	COMMON	SLANG	USE	DETECTION TIMES	PHYSICAL SYMPTOMS	DANGERS
Amphetamines (Amphetamines/ Methamphetamine)	By prescription only	Altimine Asenlix [Mex] Appetzugler [Ger] Benzedrine Captagon Desoxyn Dexedrine Didrex Eldepryl [US] Envitrol Fenistic, Esbelcaps [Mex] Geodowin Methamphetamine Methedrine Phencamine Plurimen Segontin Sydnocarb Vick's Inhalers	Benny Black beauties Bumblebees Copilots Crank Footballs Hearts Jelly Beans Meth Pep pills Speed Ups Uppers White Cross	To increase alertness and activeness. Diet Pills Stimulant	Urine: 1 to 3 days Hair: Up to 90 days (depending on length of hair) Oral Fluid: 24 to 36 hours	Anxiety Irritability Loss of appetite Mood elevation Rapid speech Tremors	Disorientation Fatigue Increase in blood pressure Paranoia Possible hallucinations Severe depression
Barbiturates	By Prescription Only	Amobarbital Butalbital Butisol Fioricet Fiorinal Phenobarbital Secobarbital	Barbs Bluebirds Blues Downers M&M's Red Devils Yellow Jackets	Sedation Seizure control	Urine: 1 to 3 days (short acting) Except Phenobarbital: 1 to 3 weeks	Decreased alertness and muscle control Depression Drowsiness Intoxication and slurred speech	Emotional instability Possible overdose and death, especially when mixed with alcohol Rigidity and painful muscle contraction

Drugs of Abuse

DRUG/FAMILY	AVAILABILITY	COMMON	SLANG	USE	DETECTION TIMES	PHYSICAL SYMPTOMS	DANGERS
Benzodiazepines	By Prescription Only	Chlorazepate Dalmane Diazepam Flurazepam Librium Medazepam Nordiazepam Oxazepam Serax Temazepam Valium Xanax	Downers	Reduce Anxiety	Urine: 1 to 14 days Oral fluid: 1 to 2 days	Disorientation Drunken behavior without odor of alcohol Slurred speech	Cold and clammy skin Coma Dilated pupils Possible death Shallow respiration Weak and rapid pulse
Cannabinoids (Marijuana)	Marinol	Marijuana	Dope Ganja Grass Mary Jane Pot Reefer Sinsemilla Smoke Weed	Psychoactive	Urine: (Occasional use): 1 to 3 days (Habitual/chronic use): Up to 30 days Hair: Up to 90 days (depending on length of hair) Oral Fluid: Less than 24 hours	Altered perception Craving for sweets Dilated pupils Increased appetite Lack of concentration and coordination Laughter	Anxiety Impaired short-term memory Increased heart rate Lung damage Possible psychosis with chronic use Psychological dependence
Cocaine & Metabolite	Cocaine, used as a vaso-constructive anesthetic in otolaryngology, ophthalmology, and dentistry	Cocaine	Big C Blow Coke Crack Flake Lady	Stimulant	Urine: 1 to 3 days Hair: Up to 90 days (depending on length of hair) Oral Fluid: 24 to 36 hours	Irritability Nervousness Short-lived euphoria changing to depression Tightness of muscles	Anxiety Fever Possible death from convulsions or respiratory arrest Shallow breathing Tremors

Drugs of Abuse

DRUG/FAMILY	AVAILABILITY	COMMON	SLANG	USE	DETECTION TIMES	PHYSICAL SYMPTOMS	DANGERS
Cocaine & Metabolite (cont.)	TAC (tetracaine, adrenaline, and cocaine mixture) used in some emergency rooms for suturing		Nose Candy Rock Roxanne Snow Snowbirds White				
Methadone	Prescription Available	Dolophine Metadose	Dolls Done	Narcotic Analgesic	Urine: 1 to 3 days	Constricted pupils Drowsiness Euphoria Nausea Respiratory depression	Clammy skin Coma Convulsions Possible death Slow and/or shallow breathing
Methaqualone	No Prescription available	Methaqualone	Lemmon Ludes Mequin Quaalude Sopor	Sedative Muscle Relaxant	Urine: 1 to 7 days	Confusion Dizziness Feeling of well being Later hangover Loss of coordination Sleepiness	Anxiety Insomnia
Opiates	Prescription available	Morphine: M Miss Emma	Codeine: -Aceta- minophen	Narcotic Analgesic	Urine: 1 to 3 days Hair: Up to 90	Euphoria Insensitivity to pain Itchiness	Hepatitis Lethargy Possible death

Drugs of Abuse

DRUG/FAMILY	AVAILABILITY	COMMON	SLANG	USE	DETECTION TIMES	PHYSICAL SYMPTOMS	DANGERS
Opiates (cont.)	Available OTC in Canada and certain other countries	Codeine: School Boy Hydrocodone: Vicodin Hydromorphone: Dilaudid Oxycodone: Percocet Percodan Oxycotin Oxymorphone: Mumorpham	with Codeine® -Actifed with Codeine Cough Syrup® -Ambenyl Cough Syrup® -Aspirin with Codeine® -Broncholate CS® -Brontex® -Calcidrine Syrup® -Capital with Codeine Suspension® -Codalan Tablets® -Codiclear DH Syrup® -Codeine phosphate®, sulfate -Codimal PH®, DH, DM, LA® -Deconsal C		days (depending on length of hair) Oral Fluid: 24 to 36 hours	Nausea Running nose Sedation Vomiting Watery eyes	Slow and shallow breathing Weight loss

Drugs of Abuse

DRUG/FAMILY	AVAILABILITY	COMMON	SLANG	USE	DETECTION TIMES	PHYSICAL SYMPTOMS	DANGERS
Opiates (cont.)			-Dimetane-DC Cough Syrup® -Empirin with Codeine® -Fiorcet with Codeine® -loTuss Liquid® -Isoclor Expectorant® - Naldecon CX Liquid® -Novahistine DH® -Nucofed® Heroin				
Phencyclidine	No Prescription available	PCP	Angel Dust Devil Stick Dummy Dust Wack	Dissociate Anesthetic [veterinary] Distortion of Senses	Urine: (Occasional use): 1 to 5 days (Habitual/chronic use): Up to 30 days Hair: Up to 90 days (depending on length of hair) Oral Fluid: 24 to 36 hours	Anxiety Mood and perception alteration Nausea Panic Paranoia Suicidal urge Tremors	Flashbacks Hallucinations Possible emotional instability and psychosis Unpredictable behavior

Drugs of Abuse

DRUG/FAMILY	AVAILABILITY	COMMON	SLANG	USE	DETECTION TIMES	PHYSICAL SYMPTOMS	DANGERS
Propoxyphene	By Prescription Only	Darvocet Darvon N Dolene		For the relief of mild to moderate pain	Urine: 1 to 3 days	Asthenia Dizziness Drowsiness Dysphoria Euphoria Headache	Convulsions Respiratory depression Skin rash and other allergic reactions occur occasionally and may be accompanied by drug fever and mucosal lesions Stupor or coma
Alcohol, Urine— Depressant liquid medications containing ethyl alcohol, as well as alcoholic beverages	OTC	Beer Liquor Wine		Sedative Depressant	Varies on metabolism	Depressant Loss of interest in activities Loss of body control / mind control Mood swings Physical dependence	Alcohol abuse Alcoholism Birth defects Cancer Heart disease Liver disease Long term health problems Pancreatitis
MDMA	No Prescription available		Adam Beans Ecstasy Hug Drug Love Drug Lover's Speed XTC X	Euphoria	Urine: 2 to 4 days Hair: Up to 90 days (depending on length of hair) Oral Fluid: 24 to 36 hours	Empathetic and unusually warm towards people Euphoria Touch and sight senses often heightened	Diaphoresis Entactogenic effects Hypertension Hyperthermia Mydriasis Stimulant Tachycardia Visual hallucinations
OxyCodone (Opiate)	By Prescription Only	OxyContin	Hillbilly Heroin	Narcotic Analgesic	Urine: 1 to 3 days	Euphoria Insensitivity to pain	Clouding of mental functions

Drugs of Abuse

DRUG/FAMILY	AVAILABILITY	COMMON	SLANG	USE	DETECTION TIMES	PHYSICAL SYMPTOMS	DANGERS
			Killers O's OC's Oceans Oxies OXY OXY's OxyCoffins OxyCotton		Oral Fluid: 24 to 36 hours	Itchiness Nausea Running nose Sedation Vomiting Watery eyes	Cold and clammy skin Coma Confusion Dizziness Loss of consciousness Nausea Reduced vision Seizures Slow breathing (respiratory depression) Small pupils Tiredness/Weakness Vomiting